

Structural Parameters of Stellar Disks from 2MASS Images of Edge-on Galaxies

Dmitry Bizyaev ^{1,2} and Sofia Mitronova ²

ABSTRACT

We present results of an analysis of the J, H, and K_s 2MASS images of 139 spiral edge-on galaxies selected from the Revised Flat Galaxies Catalog. The basic structural parameters scalelength (h), scaleheight (z_0), and central surface brightness of the stellar disks (μ_0) are determined for all selected galaxies in the NIR bands. The mean relative ratios of the scaleheights of the thin stellar disks in the J:H: K_s bands are 1.16:1.08:1.00, respectively. Comparing the scaleheights obtained from the NIR bands for the same objects, we estimate the scaleheights of the thin stellar disks corrected for the internal extinction. We find that the extinction-corrected scaleheight is, on average, 11% smaller than that in the K-band. Using the extinction-corrected structural parameters, we find that the dark-to-luminous mass ratio is, on average, 1.3 for the galaxies in our sample within the framework of a simplified galactic model. The relative thicknesses of the stellar disks z_0/h correlates with their face-on central surface brightnesses obtained from the 2MASS images. We also find that the scaleheight of the stellar disks shows no systematic growth with radius in most of our galaxies.

Subject headings: galaxies: structure, galaxies: spiral, galaxies: photometry, galaxies: halos, infrared: galaxies

1. Introduction

Study of the parameters of galactic components via decomposition of the rotation curves is often ambiguous in the case of spiral galaxies. The same rotation curve often may be explained by models with very different parameters of the components. The dark-to-light mass ratio may vary several times between the models explaining well the same rotation curve, especially if the latter is smooth and featureless (see for example de Blok et al. 2001; Bizyaev & Zasov 2002; Dutton et al. 2005). Incorporating information about the stellar velocity dispersion from observations helps to significantly constrain the masses of galactic components (Bottema 1993; Khoperskov et al. 2001) and relieve some ambiguities that appear in the modeling of rotation curves, but observing the

¹New Mexico State University and Apache Point Observatory Sunspot, NM, 88349, USA

²Sternberg Astronomical Institute, Moscow, 119992, Russia

³Special Astrophysical Observatory, Nizhniy Arkhyz, Russia

distributions of the velocity dispersion is a rather difficult task. The stellar disk thickness can be used as an alternative to the stellar velocity dispersion (Zasov et al. 2002). Studies of the vertical structure of stellar disks alone allow the constraint of some dark halo parameters (Bahcall 1984; Zasov et al. 1991, 2002; Bizyaev et al. 2003; Bizyaev & Kajsin 2004). Note that not all galaxies are suitable for such studies: inferring the structural parameters for a spiral galaxy with a significant bulge from photometric data alone is dangerous and should be supplemented with concerns of different mass-to-light ratio for different galactic subsystems (see discussion by Abadi et al. 2003, regarding difference between photometrically and dynamically identified subsystems).

Infrared observations are crucial for studies of the structure of edge-on galaxies. The dust attenuation makes the extraction of the structural parameters difficult from the optical data. In the NIR bands the galaxies are much more transparent, which facilitates studies of the structure of galaxies. The all-sky 2MASS survey (Skrutskie et al. 2006) offers a good opportunity to increase the number of edge-on galaxies available for studies in the near-infrared bands. Although faint parts of the galaxies are not seen in the 2MASS images, the thin disks of the galaxies are obtained with sufficient signal-to-noise ratio (S/N hereafter).

In this paper we focus on the determination of stellar disk structural parameters corrected for the internal extinction. One approach to the problem is to perform the modeling of the radiation transfer in 3-D stellar-gaseous disks (like in Xilouris E. et al. 1999; Kylafis et al. 2001; Bianchi 2007). Such studies require rather deep images, although there are attempts to apply this kind of analysis to larger sample of moderate quality (Bizyaev 2007). In the present paper we develop another approach to study the extinction-corrected structure of the stellar galactic disks comparing the parameters of the disks obtained from different NIR bands.

In Section 2 we discuss the sample selection and evaluation of the stellar disk’s structural parameters, consider reliability of the parameters, and study the radial variation of the disk scale-height. In Sections 3 and 4 we show how the comparison of the parameters obtained in the three NIR bands can be used to estimate the extinction-corrected scaleheight in the galactic disks. Section 5 considers the connection between the galactic disk thickness with parameters of the dark halos around the spiral galaxies in the frames of a simplified model. Section 6 summarizes results of the paper.

2. Sample of Galaxies and Analysis of the Images

As a pre-defined sample of edge-on galaxies, the Revised Flat Galaxies Catalog (RFGC hereafter, Karachentsev et al. 1999) is chosen. The catalog contains visually-classified extragalactic objects with major-to-minor axes ratio $a/b > 7$. To secure enough pixels for studying the vertical structure, we select objects with the major axis greater than 1 arcmin. This size is estimated directly from the 2MASS images at the level of $S/N \sim 3$. Note that this size is significantly less than commonly used diameter D_{25} . As a result, we select 139 spiral edge-on galaxies that have

images in all three 2MASS bands: H, J, and K_s .

We apply the technique described by Bizyaev & Mitronova (2002) to obtain the structural parameters of stellar disks of the galaxies disks. This method is based on the analysis of photometric profiles drawn parallel to the major and minor axes of galaxy at a one-pixel interval. The volume brightness in the stellar disks is assumed to change in the radial r or vertical z directions as follows:

$$\rho_L(r, z) = \rho_{L0} \exp(-r/h) \operatorname{sech}^2(z/z_0), \quad (1)$$

where h and z_0 are the scalelength and scaleheight of the disk, respectively, and $I_0 = \int \rho(0, z) dz$ is the central surface luminosity of the face-on disk, which corresponds to the central surface brightness μ_0 . The model photometric profiles are obtained by the integration of equation (1) along the line-of-sight and then by convolution with the instrumental profile. The best-fit parameters h , z_0 , and I_0 are estimated for each radial or vertical profile, and then their median values are taken as the resulting scales and central surface brightness of the stellar disks, whereas the standard deviations of h , z_0 , and I_0 estimated from different profiles are taken as uncertainty of the parameters. The central regions of the galaxies in which bulges may exist are excluded from our analysis. The parameters h , z_0 , and μ_0 are obtained in the J, H, and K_s bands for selected 139 galaxies. The K_s -based parameters are shown in Table 1.

Figure 1 shows histograms of distribution of the scale heights (top panel) and scale lengths (bottom panel) estimated for our program galaxies. In both panels, the J- and H-band scales are normalized by those in the K_s band. It can be noticed that the scaleheights look systematically larger, on average, in the J band than those in the K_s for the same galaxy. We will discuss this effect below: in § 3 using our observational data, and in § 2.2 with the help of artificial data.

2.1. Radial Variation of the Disk Scale Height

As in Bizyaev & Mitronova (2002), we investigate how the scaleheight z_0 of the stellar disks in our galaxies change with the distance to the center r . Our approach to the estimation of the z_0 is independent of the apriori assumptions about behavior of the scaleheight with radius, in a contrast with 3-D modeling of the radiation transfer in the stellar and dust disks. Since we analyze the vertical profiles separately from each other, this approach is insensitive to possible bending of the galactic disks. Note that if the resulting scaleheight $z_0(r)$ demonstrates a systematic growth with distance to the center, it suggests that the real radial gradient of the disk scaleheight may be a few times greater due to the projection effects.

We fit $z_0(r)$ with a linear function of radius r and find the radial gradients of z_0 (expressed in units of z_0) per scalelength, see Figure 2. Different lines in Figure 2 show histograms of distribution of the gradient in the J, H, and K_s bands in our whole sample. The median gradient $(dz_0/dr) \cdot (h/z_0)$ in the K_s is essentially zero (-0.01), as well as in the H (0.) and J (0.01) bands.

Note that some galaxies in histogram in Figure 2 indeed show noticeable radial gradients of the

scaleheight. In combination with projection effects, it suggests significant gradients $(dz_0/dr) \cdot (h/z_0)$ in a small fraction of our galaxies.

2.2. Reliability of the Structural Parameters

The dust disks embedded into the stellar components of the galaxies may significantly distort the structural parameters of the stellar disks estimated even from the NIR images. We check reliability of the estimated parameters via the following simulations. A set of artificial images is created using equation (1) and typical parameters h and z_0 . The surface brightness μ_0 is replaced by the luminosity surface density I_0 in our simulations. Each stellar disk contains an embedded dust disk inside with scalelength of $1.5h$ and scaleheight of $1/3 z_0$ (according to Bianchi 2007). The face-on opacity of the dust disk corresponds to $A_V = 1$ mag. This value should be close to the upper limit for the face-on extinction in spiral galaxies (Kuchinski et al. 1998; Xilouris E. et al. 1999; Bianchi 2007). Finally, the model disks are randomly inclined by 85-90 degrees and their images (i.e. 2-D projections to the plane of sky) are obtained via integration along the line-of-sight. The images are then convolved with the instrumental profile typical for the observations. Then the poisson and gaussian noise patterns are added to create sets of typically thirty artificial images with the same parameters but with different random noise pattern. The images are analyzed by the same way as the real galaxies; the mean values of h , z_0 , and I_0 and their r.m.s. are estimated and then compared with the input parameters. Figures 3 and 4 give examples how well the structural parameters can be extracted from the artificial edge-on disks made with stars and dust should they be observed in the J and K_s NIR bands. Although the scalelength and the central surface brightness can be recovered with a fairly good accuracy from both J and K_s bands, the scaleheight in the J band looks systematically thicker due to the dust absorption effects only. The K_s scaleheight in this case is found to be about 15% greater than the model input scaleheight.

We check how the chosen values of the model parameters affect our conclusions. The amount of the dust opacity is responsible for increasing the estimated scale height and makes it thicker by 10-15% in the K_s if A_V is of the order of 1 mag. Much higher values of the central dust opacity, more than 3 mag, make stellar disks look about twice as thick even in the K_s band. At the same time, the disk scalelength shows relatively small variations in this case. Variations of the dust scalelength between h and $2h$ do not change the estimated structural parameters significantly. The dust disk thickness affect the estimated stellar disk thickness in the most power when it is about a half of z_0 . Once the dust disk is getting comparable in thickness with the stellar one, it affects mostly the stellar disk surface brightness, and not z_0 .

The inclination of the stellar disk to the line-of-sight is an important parameter and it increases the stellar disk thickness estimated in our modeling by more than 15%, if the inclination is getting to be greater than 5 degrees. On the other hand, the relatively small inclination has almost no effect on the recovered stellar disk scalelength and central surface brightness. More than 5 degrees inclination in galactic disks would create problem with estimation of z_0 (see also de Grijs et al.

1997), but such galaxies should fail the RFGC catalog criteria and we don’t expect to have many of highly inclined disks in our sample. If the size of the modeling disk is so small that its true scaleheight is much less than the seeing, the resulting z_0 gets independent of the original disk scaleheight. Our selection of objects for the analysis by size of their major axis prevents such situation. Given all other parameters fixed, the recovering scalelength, scaleheight, and surface brightness are independent of the numerical value of h , z_0 , and I_0 unless z_0 is getting smaller than a pixel in the model images.

In general, our modeling suggests that we can estimate the structural parameters with of order of ten percent precision using our method. At the same time, the sensitivity to the variations of the scaleheight within the same galaxy with different amount of extinction (like in the case of observations in different NIR bands) should be by the order of magnitude better.

3. The Basic Structural Parameters in Different NIR Bands

As we can see from our observational results shown in Figure 1, the stellar disks look thicker in the J band and thinner in K_s for the same object, in agreement with our modeling from § 2.2. Two factors may be responsible for such a difference in real galaxies: the vertical gradient of stellar population properties and the dust attenuation. Let’s consider the effects of the radial and vertical gradients of stellar population in galactic disks. The radial gradients of metallicity in the thin disks are of the order of -0.07 dex/kpc for the late-type galaxies (van Zee et al. 1998), and even less in our Galaxy (Rolleston et al. 2000; Daflon & Cunha 2004; Allende Prieto et al. 2006). According to the population synthesis models by Bruzual & Charlot (2003), this creates gradients in (J- K_s) color of the order of -0.015 mag/kpc (i.e. the periphery is bluer). Hence, the scalelength in J should appear 4 % longer than that in K for a typical scalelength values of 3 kpc. A radial age gradient in combination with the metallicity change may rise this scalelength change to 5%. As we see, these numbers are within our accuracy of estimation of the scalelength. The vertical gradients of metallicity and age, and hence contribution of the stellar population effects to the vertical color variations within the thin stellar disk, are usually small (Tadross 2003; Seth et al. 2005; Allende Prieto et al. 2006). Even if we assume the large values for the vertical gradient of [Fe/H] as in de Grijs & Peletier (2000), -0.2 dex/kpc, this would create less than 1% longer scaleheights in the J than that in the K_s due to the stellar population gradients. The amplitude of the changes of scalelengths and scaleheights between the H and K_s bands is roughly a half of that between the J and K_s .

We conclude that the scaleheights of the pure thin stellar disk (i.e. without dust) estimated from the J, H, and K_s images should appear the same. All significant scaleheight variations have to be addressed to the reddening effects. This is in agreement with conclusions by Dalcanton & Bernstein (2002). Note that in some cases of extraordinary objects like a superthin galaxy UGC 7321 (Matthews & Wood 2001), the reddening alone cannot explain the observed optical color (B-R), (R-H) variations in the vertical direction and vertical gradients of metallicity

and age are required for the stellar population. At the same time, Matthews & Wood (2001) note that the superthin’s H-K color gradients might be explained by the wavelength-dependent dust absorption alone.

4. Structural Parameters of the Thin Stellar Disks Corrected for Extinction

Following Kylafis & Bahcall (1987), we consider a vertical luminosity profile of an edge-on image of a stellar disk with the volume brightness $\rho_L(z) = \rho_{L0} \exp(-z/z_0)$. Embedded co-planar dust disk with its scaleheight $z_d < z_0$ absorbs and scatters the stellar light in dependence of the dust extinction coefficient $\kappa_\lambda = \kappa_{\lambda 0} \exp(-z/z_d)$. For simplicity, we neglect the dust scatter appealing to Bianchi (2007) conclusion that the structural parameters of stellar disks in 3-D modeling of the light distribution with or without scattering are not significantly different. As in Kylafis & Bahcall (1987), the brightness distribution along the minor photometric axis (z) in such a disk is $I(z) = I_0(z) \frac{1 - \exp(-\tau(z))}{\tau(z)}$, where $I_0(z)$ is the brightness profile in the case of zero extinction, and τ is the optical depth across the whole edge-on disk, which also depends on the vertical distance over the galactic plane z . For small τ , $I(z) \approx I_0(1 - 0.5\tau(z))$. The observed scaleheight z_e of the stellar disk with added extinction is connected with its projected brightness along the selected profile as $I(z) = I_{ce} \exp(-z/z_e)$. Given $I_0(z) = I_c \exp(-z/z_s)$, we find that $1/z_e \approx 1/z_s(1 - \tau(0))$. Therefore, estimating the thickness of the stellar disk with dust from its edge-on profiles we expect to find

$$z_{e\lambda} = z_s(1 + C_\lambda \tau_V), \quad (2)$$

where $z_{e\lambda}$ is the z_e estimated from the disk’s brightness profile at a certain photometric band, and the optical depth in the plane of the disk in certain photometric band is $\tau_\lambda = C_\lambda \tau_V$. We assume that C_λ equals to 0.276, 0.176, and 0.112 in the J, H, and K_s bands, respectively (Schlegel et al. 1998). Since the V-band optical depth τ_V and z_s are fixed for the same disk, we use linear regression and fit a line to three data points (for J, H, and K_s values) at $C_\lambda - z_{e\lambda}$ diagram to find z_s .

For the convenience of comparison of the vertical scales, we normalize the J and H scaleheights in each galaxy by their K_s -band scaleheight. The upper panel in Figure 5 compares the normalized scaleheights. Each line in the panel connects three thicknesses for one galaxy. The tendency to observe the stellar disks thicker in the J than in the K_s is well seen. The histogram in the lower panel in Figure 5 shows the distributions of all normalized z_0 in three NIR colors. The vertical thick line designates all scaleheights in the K_s because of the normalization. The mean ratio of z_0 in the J and H to that in K_s is 1.16 and 1.08, respectively.

We calculate the scaleheight z_s corrected for the extinction (or extinction-free hereafter) using equation (2), which gives $z_s \leq z_0(K_s)$. The distribution of z_s normalized by the $z_0(K_s)$ is also shown in the lower panel in Figure 5. The median $z_s/z_0(K_s)$ is about 0.89, i.e. the thin stellar disks in our galaxies are 11% thinner than it can be estimated from the analysis of the vertical profiles from the K_s -band images. The extinction-free scaleheights are given in Table 1.

We observe no systematic variations in the scalelength between the NIR bands, as it can be seen in Figure 6, which is the same as Figure 5 but drawn for the scalelengths. This is not a surprise because our simulations shown in Figures 3 and 4 suggest rather small systematic variations of the radial scalelength among the NIR bands, and proximity of the h estimated from the J, H, and K_s bands to its extinction-corrected value. Using the scalelength h from the K_s band and extinction corrected scaleheights we find that the mean radial-to-vertical scale ratio in the thin extinction-free stellar disks of galaxies in our sample is 5.6.

5. Thickness of the Stellar Disks and Mass of the Dark Halos

The thickness of the stellar disk is sensitive to the gravitational potential of the disk + bulge + halo system in the vertical direction (e.g. Bahcall 1984). Therefore, the relative thickness of the stellar disk z_0/h may provide some information about the spherical-to-disk mass ratio. As a first step, we consider a very simplified model of a galaxy that consists of a stellar exponential disk embedded into a spherical dark halo.

The total mass of the exponential stellar disk with the central surface density Σ_0 is

$$M_d = 2\pi\Sigma_0 h^2. \quad (3)$$

The total mass of all components of the galaxy is

$$M_t \sim V^2 h, \quad (4)$$

where V is the disk’s rotational velocity. Following Zasov et al. (1991) and Kregel et al. (2005), we assume that the disk thickness depends on the local surface density Σ and vertical velocity dispersion σ_z :

$$z_0 \sim \sigma_z^2 / G\Sigma \quad (5)$$

and the ratio of the vertical-to-radial velocity dispersions is $\sigma_z/\sigma_r = \text{const}$. This is a rather coarse but fair approximation, see Polyachnko & Shukhman (1977); Bottema (1993); Gerssen et al. (2000); Kregel et al. (2005).

We incorporate an assumption of the marginal stability of the stellar disk, i.e.

$$\sigma_r = Q \cdot 3.36\Sigma/\varkappa, \quad (6)$$

where the Toomre parameter $Q = \text{const}$ (Toomre 1964). Here $\varkappa$ designates the epicyclic frequency, and $\varkappa \sim V/h$ for the flat rotation curves (see Binney & Tremaine 1987, p.121). Substituting σ_z and Σ from equations (3), (4), and (6) into (5), we find $z_0 \sim Q^2 \Sigma^2 / (\varkappa^2 G \Sigma) \sim \Sigma h^2 / (V^2 G) \sim (M_d/M_t)h$. Therefore,

$$z_0/h \sim (M_d/M_t). \quad (7)$$

Numerical N-body simulations (Zasov et al. 1991; Mikhailova et al. 2000) in which the spherical subsystem was introduced as a fixed potential confirm the equation (7). Figure 7 is adopted from Mikhailova et al. (2000) with additions from later simulations by Khoperskov (private communication). Figure 7 shows that the stellar disk thickness z_0/h is sensitive to the relative mass of the spherical component M_s normalized by the M_d . Both M_s and M_d are determined within the limits of the stellar disk (four scalelengths in the simulations). Circles in Figure 7 correspond to different numerical models with various parameters of the disk and dark halo subsystems, as well as with different bulges. The upper curve corresponds to the bulgeless models, whereas the lower one is for the models with some bulge contribution.

Contemporary and more realistic N-body simulations include "live" halos that respond to the gravitational potential of disks. The relation z_0/h versus M_s/M_d should look very similar in those simulations as in Figure 7 for the galaxies in which the spherical subsystems dominate by mass (i.e. for large M_s/M_d). For the spherical subsystems with masses $M_s \lesssim M_d$, Figure 7 illustrates an extreme case, and its top left part (large z_0/h and small M_s/M_d) demonstrates the lower limit for mass of the spherical subsystem given disk thickness z_0/h . Preliminary results from N-body simulations (Khoperskov, private communication) with live halo suggest not essential corrections to M_s/M_d inferred from z_0/h according to Figure 7 (of the order of 15%). We encourage reader to assume our further estimations of halo-to-disk mass ratio made from the disk thickness as lower limits for the real M_s/M_d .

Following Figure 7, we calculate the distribution of the dark halo mass M_{halo} expressed in the units of the stellar disk mass for our galaxies. The RFGC members are "flat" galaxies that are not expected to harbor large bulges, so we assume here that $M_{halo} = M_s$. We find the following relation for bulgeless galaxies from Figure 7: $M_{halo}/M_d = -0.9419 + 0.3737(h/z_0)$. According to this formula, the median value of M_{halo}/M_d is about 1.3 within our sample. The histogram of distribution of M_{halo}/M_d is shown in Figure 8. No any correlations between M_{halo}/M_d and absolute magnitude, maximum of rotational velocity, or radial gradient of the scaleheight is found for our galaxies. A tendency to observe higher M_{halo}/M_d in the objects with larger scalelength h is found, although this relation shows large uncertainty.

Following the same basic assumptions, we can include the central surface density Σ_0 into consideration. As it follows from our definition of the disk and total galaxy mass, $M_d/M_t \sim \Sigma_0 h^2/V^2 h = \Sigma_0 h/V^2$. We perform a linear fit between $\log V$ and $\log h$ for our galaxies with known V and found that $h \sim V^{1.42}$. It seems more correct to utilize non edge-on galaxies to study such a correlation, and Kormendy (1990) and Walker (1999) suggest $h \sim V^{1.4}$ and $h \sim V^{1.5}$, respectively, for arbitrary inclined disks. If the latter relation is accepted, it gives $M_d/M_t \sim \Sigma_0/V^{0.5}$, and in combination with equation (7) we come to the following relation between the disk thickness and its central surface density

$$z_0/h \sim \Sigma_0/h^{1/3}. \quad (8)$$

The upper panel in Figure 9 shows the relation between h/z_s (i.e. extinction-corrected inverse thickness of the disks) and the central surface brightness in the K_s band $\mu_0(K)$. Here $\mu_0(K)$ was

corrected for the extinction according to formula (2). The correction is rather small, and its median value for our sample of galaxies is about 0.1 mag/arcsec².

The lower panel in Figure 9 demonstrates the plot of h/z_s versus the central surface density estimated from observations as $2.30(M/L)2.512^{24-\mu_0}$. The numerical coefficient in the formula corresponds to the K_s -band solar absolute magnitude 3.33 mag. We assume the constant $(M/L_{K_s}) = 1$ here. The trend in Figure 9 is in agreement with formula (8). The very big scatter of points in Figure 9 can be attributed to the relatively low accuracy in the μ_0 , h , and z_s , and even more uncertainty is due to the non-constant value of the mass-to-light ratio (M/L) . The scatter of this relation can be lowered dramatically if only large ($h > 4$ kpc) galaxies with the scaleheight gradient in the K_s $(dz_0/z_0)/(dr/h) < 0.05$ are considered, see Figure 10. In this case the inverse disk thickness (h/z_s) depends on $\mu_0(K)$ as

$$h/z_0 = 47.557 - 23.069x + 3.016x^2, \quad (9)$$

where $x = \log(2.3 \cdot 2.512^{24-\mu_0(K)}) = 0.4 * (25.11 - \mu_0(K))$. The solid curve in Figure 10 represents this equation. Apparently, the large spiral galaxies whose stellar disks are not flared can be better described by the toy-model developed above. Establishing the correlation between the μ_0 and h/z_s may be useful for estimating the disk thickness and M_{halo}/M_d in disk galaxies with arbitrary inclination.

Note that our model is oversimplified, and attempts to estimate the M_{halo}/M_d ratio only. Should we need to know the values of M_{halo} or M_d taken apart, or structural parameters of the dark halo, the rotation curve modeling has to be incorporated. Such additional parameter as the disk thickness helps to decrease the ambiguity during the rotation curves decomposition. A special attention should be given to correspondence between the photometrically and dynamically inferred parameters of galactic subsystems. An example of N-body simulations of an early-type disk galaxy (Abadi et al. 2003) demonstrates that photometry may fail to trace the dynamically distinctive subsystems in the central parts of bulge-harboring galaxies. In our consideration we avoid central parts of galaxies, and using the RFGC objects prevents us from considering bulge-dominated stellar systems.

The weakness of the simplified assumptions in the modeling described above encourages us to apply a more realistic simulations of the stellar disk vertical structure. Since it requires extensive numerical simulations and better quality infrared data, and will gain from the incorporation of rotation curves decomposition, we defer such more complex consideration to a further paper.

6. Summary and Conclusions

We study how the dust absorption affects the basic structural parameters (disk central surface brightness, vertical and radial scales) of stellar disks in spiral galaxies estimated from the NIR J,H, and K_s images. Using 2MASS data, we compare the structural parameters estimated from

different NIR bands focusing on the scaleheight of thin stellar disks. The stellar disks look thinner in the 2MASS K_s band in the comparison with the H and J. We employ this fact to figure out the extinction-corrected scaleheight z_0 of the thin stellar disk.

Using 139 relatively large galaxies selected from the 2MASS catalog, we find that the mean vertical scaleheight (z_0) ratios are 1.16:1.08:1.00:0.89 in the J:H: K_s :extinction-free bands, respectively. This means that the radial scalelength estimated from the K_s images is very close to the extinction-corrected one, whereas the scaleheight is overestimated by 11% on average. The mean extinction correction for the face-on central surface brightness is only about 0.1 mag/arcsec² in the K_s .

The median radial-to-vertical scale ratio for our sample is about 6. Using a relation between the stellar disk thickness and the halo-to-disk mass ratio obtained from a simplified model, we estimate the dark-to-luminous mass ratio in our galaxies within the limits of their optical disks. Its mean value is about 1.3. A relation between the stellar disk thickness and its central surface brightness is observed for the galaxies from our sample, although it shows significant scatter and is affected by uncertainty in additional parameters. This relation is more prominent for large ($h > 4$ kpc) galaxies from our sample and can be utilized for coarse estimating the disk thickness and relative mass of the dark halo in large spiral galaxies arbitrarily inclined to the line-of-sight.

We find also that the scaleheight does not change systematically along the radius in most of our galaxies. Only a small fraction of galaxies demonstrates a noticeable radial growth of the scaleheight.

This project was partly supported by grant RFBR-07-02-00792. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. Computational equipment for DB was provided by AAS under a Small Research Grant. We appreciate recommendations from anonymous referee that improved the paper. We acknowledge the usage of the HyperLeda database (<http://leda.univ-lyon1.fr>).

REFERENCES

- Abadi M., Navarro J., Steinmetz M., & Eke V. 2003, *ApJ*, 597, 21
- Allende Prieto, C., Beers, T. C., Wilhelm, R., et al. 2006, *ApJ* 636, 804
- Bahcall, J. 1984, *ApJ*, 276, 156
- Bianchi, S. 2007, *A&A*, 471, 765
- Binney, J. & Tremaine, S., 1997, "Galactic dynamics", Princeton, NJ, Princeton University Press

- Bizyaev, D. 2000, *Astronomy Letters* 26, 219
- Bizyaev, D., Mitronova, S. 2002, *A&A*, 389, 795
- Bizyaev, D., Zasov, A., 2002, *Astronomy Reports*, 46, 721
- Bizyaev, D., Khoperskov, A., & Tiurina, N. 2003, *Bull. Am. Astron. Soc.*, 35, 758 (202.4013)
- Bizyaev, D. & Kajsin, S. 2004, *ApJ*, 613, 886
- Bizyaev, D. 2007, *Bull. Am. Astron. Soc.*, 39, 758 (211.1323)
- de Blok W., McGaugh S., & Rubin V. 2001, *AJ*, 122, 2396
- Bottema, R. 2001, *A&A*, 275, 16
- Bruzual, G. & Charlot, S. 2003, *MNRAS*, 344, 1000
- Daffon, S. & Cunha, K. 2004, *ApJ*, 617, 1115
- Dalcanton, J. & Bernstein, R. 2002, *AJ*, 124, 1328
- Dutton, A., Courteau, S., de Jong, R., et al. 2005, *ApJ*, 619, 218
- Gerssen, J., Kuijken, K., & Merrifield, M. 2000, *MNRAS*, 317, 545
- de Grijs, R. & Peletier, R. 2000, *MNRAS*, 313, 800
- de Grijs, R., Peletier, R. F., van der Kruit, P. C. 1997, *A&A*, 327, 966
- Karachentsev, I. D., Karachentseva, V. E., Kudrya, Yu. N., et al., 1999, *Bull. Spec. Astr. Obs.*, 47, 5
- Khoperskov A.V., Zasov, A.V., & Tiurina, N. 2001, *Astron. Rep.*, 45, 180
- Kormendy, J. 1990, in "Evolution of the universe of galaxies", proc. of the Edwin Hubble Centennial Symposium, Berkeley, CA, June 21-23, p.33
- Kregel, M., van der Kruit, P. C., Freeman, K. C. 2005, *MNRAS*, 358, 503
- Kregel, M. & van der Kruit, P., *MNRAS* 2005, 481, 358
- Kuchinski, L., Terndrup, D., Gordon, K. et al. 1998, *AJ*, 115, 1438
- Kylafis, N., & Bahcall, J. 1987, *ApJ*, 317, 637
- Kylafis, N., Misiriotis, A., Papamastorakis, J., & Xilouris, E. 2001, *Astrophys. and Space. Sci.* 276, 531
- Matthews L. & Wood, K. 2001, *ApJ*, 548, 150

- Mikhailova, E., Khoperskov, A., et al. 2000, in "Stellar Dynamics: From Classic to Modern", Proc. conf. Saint Petersburg, August 21-27, 2000
- Polyachnko, V. L. & Shukhman, I. G. 1977, *Sov. Astronomy Letters*, 3, 134
- Rolleston, W. R. J., Smartt, S. J., Dufton, P. L., et al. 2000, *A&A*, 363, 537
- Seth, A., Dalcanton, J., & de Jong, R. 2005, *AJ*, 130, 1574
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Skrutskie, M.F., Cutri, R.M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Tadross, A. L. 2003, *New Astronomy*, 8, 737
- Toomre, A. 1964, *ApJ*, 139, 1217
- van Zee, L., Salzer, J., Haynes, M., et al. 1998, *AJ*116, 2805
- Walker, M. 1999, *MNRAS*, 308, 551
- Xilouris, E.M., Byun, Y.I., Kylafis, N.D. et al. 1999, *A&A*, 344, 868
- Zasov, A.V., Makarov, D., & Mikhailova, E. 1991, *Sov. Astron. Lett.*, 17, 374
- Zasov, A., Bizyaev, D., Makarov, D., Tiurina, N. 2002, *Astron. Letters*, 28, 527

Table 1: The Structural Parameters of the Galaxies

Name	$h(K)$	dh	$z_0(K)$	dz_0	$\mu_0(K)$	$d\mu_0$	$z_0(\text{cor.})$	$z_0(\text{cor.})/h$	$\mu_0(K, \text{cor.})$	$\frac{M_{\text{dark}}}{M_{\text{lum}}}$
	kpc	kpc	kpc	kpc	mag/□"	mag/□"	kpc	–	mag/□"	–
RFGC0095	0.72	0.06	0.31	0.08	17.69	0.26	0.27	0.375	17.61	0.06
RFGC0099	4.35	0.06	0.95	0.17	17.51	0.18	0.83	0.191	17.41	1.02
RFGC0102	1.76	0.03	0.48	0.11	17.49	0.20	0.46	0.260	17.46	0.49
RFGC0124	8.15	0.24	1.36	0.19	17.94	0.18	1.12	0.137	17.81	1.79
RFGC0152	5.33	0.06	0.88	0.14	17.73	0.21	0.76	0.143	17.66	1.67
...										
The full table is available on-line										

The table shows RFGC name, scalelength h in kpc and its uncertainty in the K_s band, scaleheight z_0 in kpc and its uncertainty in the K_s , central surface brightness μ_0 in the K_s , extinction-corrected scaleheight $z_0(\text{cor.})$ in kpc, extinction-corrected relative thickness of the stellar disk $z_0(\text{cor.})/h$, extinction-corrected central surface brightness $\mu_0(\text{cor.})$ in the K_s , and dark-to-luminous mass ratio $M_{\text{dark}}/M_{\text{lum}}$.




Fig. 1.— The scaleheights (top panel) and scalelengths (bottom panel) in the J (dashed line) and H (dotted line) bands normalized by those from the K_s for our 139 galaxies. The K_s band scales in both panels equal to unit for all objects.




Fig. 2.— Distribution of the radial gradients of the scaleheight per one scalelengths $(dz_0/z_0)/(dr/h)$. The solid, dashed, and dash-dotted lines designate the K_s , H, and J gradients, respectively. The three short vertical lines above the histogram indicate the median values of the gradients in the K_s , H, and J bands (from left to right), that are essentially equal to zero.




Fig. 3.— The structural parameters of the dusty stellar disks estimated from sets of artificial images in the J band are shown by the solid lines with squares for different signal-to-noise ratio (S/N in the center of the disks is used). The dotted lines designate the r.m.s scatter of the parameters. The dash-dotted horizontal lines denote the input (i.e. true) model parameters. The scalelength, scaleheight, and central surface luminosity (I_0 , in arbitrary units) are shown in the upper, middle, and lower panels, respectively.




Fig. 4.— The same as in Figure 3 but for the K_s band.




Fig. 5.— Top panel: all scaleheights z_0 normalized by that in the K_s . Each a galaxy is shown by a dashed line, which connects the z_0 values in the J, H, and K_s (the latter equals to unit for all objects). The solid line shows the case of equality of all scaleheights. Bottom panel: histograms of distribution of the scaleheights normalized by those in the K_s band. Dashed line corresponds to the J band, dotted - to the H band. The solid line designates the extinction-corrected scaleheights. The numbers under the letters show the mean values of the scaleheights in the corresponding bands. The thick vertical line designates formally the scaleheights in the K_s that are all equal to unity due to the normalization.




Fig. 6.— Top panel is the same as in Figure 5 but drawn for the disk scalelengths. Bottom panel is similar to the bottom panel in Figure 5 and is plotted for the scalelengths. The scalelengths are normalized by those from the K_s band. Dashed line corresponds to the J band, dotted - to the H band. The thick vertical line designates formally the scalelengths in the K_s that are all equal to unity due to the normalization.




Fig. 7.— The relative disk thickness z_0/h is shown against the spherical-to-disk mass ratio M_s/M_d . Here M_s is the mass of the spherical component (halo and bulge, in a general case), and M_d is the mass of the disk component. The M_s and M_d are estimated within the limits of the stellar disk ($4h$). This figure comes from simplified N-body simulations (Mikhailova et al. 2000) and each symbol corresponds to certain numerical model. The models were run with a variety of parameters for the disk, dark halo, and bulge. The upper curve corresponds to the bulgeless models, whereas the lower one is for the models with some bulge contribution.




Fig. 8.— The dark halo-to-disk mass ratio for the galaxies from our sample inferred from the relative disk thickness. The mass ratio is close to the dark-to-luminous mass ratio since most of our galaxies have no prominent bulges.




Fig. 9.— Top panel: the inverse disk thickness h/z_s corrected for the internal extinction versus the central surface brightness in the K_s band. Bottom panel: the same relation where μ_0 is replaced with the central surface density calculated as $2.30 \cdot 2.512^{24 - \mu_0}$, see text.




Fig. 10.— The inverse disk thickness h/z_s corrected for the extinction versus the central surface density of the stellar disks (see text) for the sample of large ($h > 4$ kpc) and non-flared disk galaxies. The solid line shows the best-fit to the data (see text).